



FOCINVEZ™ is a prescription medicine used in patients 6 months of age and older to prevent nausea and vomiting caused by certain anti-cancer medicines (chemotherapy). When used for this purpose, FOCINVEZ™ is always used with other medicines that treat nausea and vomiting.

FOCINVEZ™ is not used to treat nausea and vomiting that you already have. It is not known if FOCINVEZ™ is safe and effective in children less than 6 months of age.

## Discover how FOCINVEZ™ can help



Look inside  
to find:



What FOCINVEZ™ is



What to expect when  
taking FOCINVEZ™



How FOCINVEZ™  
is given



Resources  
for support

Amneal  
**PATHways**  
Patient Access To Health

### IMPORTANT SAFETY INFORMATION

Do not take FOCINVEZ™ if you are allergic to fosaprepitant, aprepitant, or any of the ingredients in FOCINVEZ™.  
Do not take FOCINVEZ™ if you are taking pimozide (ORAP).

Please see Important Safety Information on pages 6-7 and Prescribing Information available in the pocket or online at [Focinvez.us](http://Focinvez.us).



Biosciences

# What is FOCINVEZ™?

FOCINVEZ™ is a prescription medicine that helps to prevent nausea and vomiting caused by some types of chemotherapy



**FOCINVEZ™ doesn't treat cancer, but it is a part of some cancer treatment plans.** Some kinds of chemotherapy that treat cancer are more likely to cause side effects of nausea and vomiting. FOCINVEZ™ is given before these kinds of chemotherapy to help prevent nausea or vomiting. When used for this purpose, FOCINVEZ™ is always used with other medicines that treat nausea and vomiting. FOCINVEZ™ is not used to treat nausea and vomiting that you already have.



**FOCINVEZ™ is approved for use in patients 6 months of age and older.** It is not known whether FOCINVEZ™ is safe and effective in children less than 6 months of age.

*If you have any questions about FOCINVEZ™, ask your healthcare provider.*


## IMPORTANT SAFETY INFORMATION

Before you take or receive FOCINVEZ™, tell your doctor about all of your medical conditions, including if you:

- Have liver problems
- Are pregnant or plan to become pregnant
- Are breastfeeding or plan to breastfeed

Please see Important Safety Information on pages 6-7 and Prescribing Information available in the pocket or online at [Focinvez.us](http://Focinvez.us).

# How is FOCINVEZ™ given?



**For adults 18 years of age and older, FOCINVEZ™ is given by intravenous (IV) infusion in the patient's vein**

- |                  |   |
|------------------|---|
| <b>WHEN?</b>     | ▶ Day 1 of chemotherapy                     |
| <b>DURATION?</b> | ▶ 20 to 30 minutes                          |
| <b>Timing?</b>   | ▶ Completed ~30 minutes before chemotherapy |

**For children 6 months to 17 years of age, FOCINVEZ™ is given through a type of IV called a central line**

- |                  |   |
|------------------|---|
| <b>WHEN?</b>     | ▶ Day 1 of chemotherapy*                    |
| <b>DURATION?</b> | ▶ 30 to 60 minutes†                         |
| <b>Timing?</b>   | ▶ Completed ~30 minutes before chemotherapy |

*If you take the blood thinner warfarin (also known as Coumadin® or Jantoven®), your healthcare provider may do blood tests to check your blood clotting after you receive FOCINVEZ™.*

\*Depending on your child's chemotherapy treatment plan, they may also take aprepitant by mouth in a tablet or liquid form on Days 2 and 3. If your child will receive oral aprepitant, see the Patient Information for those products for further information.

†The duration of infusion for FOCINVEZ™ is 30 minutes for patients 12 years to 17 years of age and 60 minutes for patients 6 months to less than 12 years of age.



# What are the possible side effects of FOCINVEZ™?

Side effects can occur during treatment with FOCINVEZ™. It is important to stay in close communication with your or your child's healthcare provider throughout your treatment.

## Serious side effects can occur with FOCINVEZ™. Contact your healthcare provider right away if you experience any of the following:

- Signs of a serious allergic reaction, such as:
  - Hives, rash, itching, flushing or redness of your face or skin
  - Trouble breathing or swallowing
  - Dizziness
  - A rapid or weak heartbeat
  - Feeling faint during or soon after you receive FOCINVEZ™
- Signs of severe skin reactions, such as rash, skin peeling, or sores
- Signs of an infusion site reaction, such as pain, swelling, and redness where the IV was placed in your or your child's body

## In adults, the most common side effects of FOCINVEZ™ include:

- Tiredness
- Diarrhea
- Low white blood cell and red blood cell counts
- Weakness
- Feeling weak or numb in your arms and legs
- Painful or difficult digestion, or changes in digestion (dyspepsia)
- Urinary tract infection
- Pain in your arms and legs

## In children 6 months to 17 years of age, the most common side effects of FOCINVEZ™ include:

- Low red blood cell count
- Low white blood cell count
- Low blood platelet count
- Low white blood cell count with a fever

Please see Important Safety Information on pages 6-7 and Prescribing Information available in the pocket or online at [Focinvez.us](http://Focinvez.us).

# Resources for support

## Learning more about cancer

You may find it useful to know more about your or your child's diagnosis and treatment. **Below are some websites that could help you understand what to expect during cancer treatment.**

American Cancer Society	CDC	The ASCO Foundation	National Cancer Institute
<a href="http://www.cancer.org">www.cancer.org</a>	<a href="http://www.cdc.gov">www.cdc.gov</a>	<a href="http://www.conquer.org">www.conquer.org</a>	<a href="http://www.cancer.gov">www.cancer.gov</a>

## The Amneal PATHways® program is here to make accessing and paying for medicine as seamless as possible

**When you enroll in the program, you will be connected with one of our Patient Access Specialists who will:**

- Help you understand your benefits and insurance coverage
- Discuss affordability options you may be eligible for
- Keep you informed about program updates as they occur

### The Amneal PATHways® Co-pay Assistance Card



Eligible\* commercially insured patients may pay as little as \$0 per treatment with an Amneal product.



Scan the QR code to enroll in the Amneal PATHways® program, and see what affordability options you may be eligible for.

Questions? Call **1-866-4AMNEAL (1-866-426-6325)** with any questions or concerns

Monday-Friday, 8:00AM-8:00PM ET

\*Eligibility conditions apply. Please contact your Patient Access Specialist for more information.



## Approved Use

FOCINVEZ™ is a prescription medicine used in patients 6 months of age and older to prevent nausea and vomiting caused by certain anti-cancer medicines (chemotherapy). When used for this purpose, FOCINVEZ™ is always used with other medicines that treat nausea and vomiting.

FOCINVEZ™ is not used to treat nausea and vomiting that you already have.

It is not known if FOCINVEZ™ is safe and effective in children less than 6 months of age.



Please see Prescribing Information available in the pocket or online at [Focinvez.us](http://Focinvez.us).

## IMPORTANT SAFETY INFORMATION

**Do not take FOCINVEZ™ if you are allergic to fosaprepitant, aprepitant, or any of the ingredients in FOCINVEZ™.**

**Do not take FOCINVEZ™ if you are taking pimozide (ORAP).**

**FOCINVEZ™ may cause serious side effects, including:**

- **Serious allergic reactions, such as anaphylaxis. Get emergency medical help right away if you get any of the following symptoms of a serious allergic reaction:** hives, rash, itching, flushing or redness of your face or skin, trouble breathing or swallowing, dizziness, a rapid or weak heartbeat, or if you feel faint during or soon after you receive FOCINVEZ™. Severe skin reactions, which may include rash, skin peeling, or sores, may occur.
- **Infusion site reactions (ISRs) at or near the infusion site have happened with FOCINVEZ™.** Most severe ISRs have happened with a certain type of chemotherapy medicine that can burn or blister your skin (vesicant) with side effects, including pain, swelling, and redness. Death of skin tissue (necrosis) has happened in some people getting this type of chemotherapy medicine. Most ISRs can happen with the first, second, or third dose and some can last up to 2 weeks or longer. Tell your healthcare provider right away if you get any infusion site reactions.

**In adults, the most common side effects of FOCINVEZ™ include** tiredness; diarrhea; low white blood cell and red blood cell counts; weakness; feeling weak or numb in your arms and legs; painful, difficult, or changes in your digestion (dyspepsia); urinary tract infection; and pain in your arms and legs.

**In children 6 months to 17 years of age, the most common side effects of FOCINVEZ™ include** low red blood cell count, low white blood cell count, low blood platelet count, and low white blood cell count with a fever.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of FOCINVEZ™. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects.

**Before you take or receive FOCINVEZ™, tell your doctor about all of your medical conditions, including if you:**

- **Have liver problems**
- **Are pregnant or plan to become pregnant.** It is not known if FOCINVEZ™ can harm your unborn baby. Women who use birth control medicines containing hormones to prevent pregnancy (birth control pills, skin patches, implants, and certain IUDs) should also use a backup method of birth control that does not contain hormones, such as condoms and spermicides, during treatment with FOCINVEZ™ and for 1 month after receiving the last dose of FOCINVEZ™.
- **Are breastfeeding or plan to breastfeed.** It is not known if FOCINVEZ™ passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you receive FOCINVEZ™.

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements. FOCINVEZ™ and certain other medicines can affect each other, causing serious side effects.

**For more information about all the side effects of FOCINVEZ™, please read the Patient Information located at the end of the Prescribing Information. You can also read the full Prescribing Information that is written for health professionals.**

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.



# Supportive care for cancer treatment



**We at Amneal Biosciences are here to support you with every aspect of the treatment journey**

Treating cancer is a team effort. Our trusted patient support services partner with the healthcare provider and treatment team, so patients can start treatment with FOCINVEZ™ knowing there will always be someone there to help.



**If you have any questions about FOCINVEZ™, ask your healthcare provider.**



**Learn about affordability options offered through the PATHways® patient support program**



## **IMPORTANT SAFETY INFORMATION**

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. FOCINVEZ™ and certain other medicines can affect each other, causing serious side effects.

**Please see Important Safety Information on pages 6-7 and Prescribing Information available in the pocket or online at [Focinvez.us](https://www.focinvez.us).**

**[amnealbiosciences.com](https://www.amnealbiosciences.com) | [amneal.com](https://www.amneal.com)**

FOCINVEZ™ is a trademark of Amneal Pharmaceuticals LLC.  
Coumadin® is a registered trademark of Bristol-Myers Squibb Company.  
Jantoven® is a registered trademark of Upsher-Smith Laboratories, LLC.  
©2024 Amneal Pharmaceuticals LLC. All rights reserved. PP-PAT-FOC-US-0001 06/24

